

CLEAN Meds Clinician Scientist Meeting Decisions  
 July 2017

| <b>Suggested Change</b>                             | <b>Collaborative Recommendation</b> | <b>Collaborative GRADE</b> | <b>Final Decision</b>                                       |
|---|-------------------------------------|----------------------------|---|
| Formoterol should be added to replace salmeterol    | Strongly against addition           | High                       | Formoterol was not added as a replacement for salmeterol    |
| Terazosin should be added                           | Strongly against addition           | High                       | Terazosin was not added                                     |
| Dutasteride should be added                         | Strongly against addition           | High                       | Dutasteride was not added                                   |
| Citalopram should be added                          | Strongly against addition           | High                       | Citalopram was not added                                    |
| Buspirone should be added                           | Strongly against addition           | High                       | Buspirone was not added                                     |
| Antibiotics for conjunctivitis should be added      | Strongly for addition               | High                       | Antibiotics for conjunctivitis was added                    |
| Aripiprazole should be added to replace risperidone | Strongly against addition           | High                       | Aripiprazole was not added as a replacement for risperidone |
| Ezetimibe should be added                           | Weakly against addition             | Low                        | Ezetimibe was not added                                     |
| Liraglutide should be added                         | Weakly against addition             | Low                        | Liraglutide was not added                                   |
| Empagliflozin should be added                       | Weakly against addition             | Low                        | Empagliflozin was not added                                 |
| Erythropoietin should be added                      | Weakly against addition             | Low                        | Erythropoietin was not added                                |